

Spinons as composite fermions

Daniel C. Cabra¹ and Gerardo L. Rossini²

*Departamento de Física,
Universidad Nacional de La Plata,
C.C. 67 (1900) La Plata, Argentina.*

ABSTRACT

We show that gauge invariant composites in the fermionic realization of $SU(N)_1$ conformal field theory explicitly exhibit the holomorphic factorization of the corresponding WZW primaries. In the $SU(2)_1$ case we show that the holomorphic sector realizes the spinon $Y(sl_2)$ algebra, thus allowing the classification of the chiral Fock space in terms of semionic quasi-particle excitations created by the composite fermions.

¹CONICET, Argentina. E-mail address: cabra@venus.fisica.unlp.edu.ar

²CONICET, Argentina. E-mail address: rossini@venus.fisica.unlp.edu.ar

i) Introduction

The notion of generalized statistics has been the subject of active research in the last ten years. Firstly, it was proven that particles obeying fractional statistics, intermediate between bosons and fermions, can exist in $2 + 1$ dimensions [1]. Then, the possibility that also $1 + 1$ dimensional systems admit fractional statistics excitations was recognized (in this case the notion of generalized statistics must be introduced) [2, 3]. Now, in spite of the amount of effort devoted to the subject, a general formulation of quantum statistical mechanics for systems whose excitations obey fractional statistics, as spinons in $1 + 1$ or anyons in $2 + 1$ dimensions, has not been fully developed, although progress in this direction has been made in Refs. [3, 4].

Concerning the physical interest of these matters, a physical manifestation of $1 + 1$ fractional statistics behavior in continuous systems has arisen, for instance, in the study of the incompressible quantum Hall liquid. It appears in connection with the two-dimensional modes in the boundary whose dynamics is governed by a Conformal Field Theory (CFT) [5]. The study of field theories whose excitations obey generalized statistics has also received much attention in connection with the so-called spinon representation of CFT [2, 6, 7, 8].

For the reasons above, it is important to have as much information as possible concerning $1 + 1$ dimensional systems manifesting these properties, particularly that arising from an explicit Lagrangian formulation (which is lacking). A thorough comprehension of the system properties, including its statistical and thermodynamical behavior, could be better achieved in this context.

The purpose of the present note is twofold:

i) We show that the $SU(N)_1$ WZW primary fields can be *explicitly* factorized into holomorphic and anti-holomorphic fields (a fact that is usually heuristically assumed). The holomorphic factors correspond to the simplest BRST invariant fields in a fermionic coset representation of the model.

ii) In connection with the recently revealed Yangian structure of these theories [6, 7, 8, 9] we show that the BRST invariant fields mentioned above exactly behave as the spinon fields of Refs. [7, 8]. In this way we provide a fermionic *Lagrangian* formulation for spinons, in the form of a well-known two-dimensional field theory in which the physical (BRST invariant) operators naturally exhibit $1 + 1$ generalized statistics [3].

We start by studying the $SU(N)_1$ -WZW CFT formulated as a fermionic coset model. This representation provides an explicit holomorphic factorization of the primary fields, in which the holomorphic factors are indeed given by the BRST invariant composite fermions.

As discussed in Refs. [6, 7, 8, 9], there exists an underlying Yangian $Y(sl_N)$ symmetry in the $SU(N)_1$ -WZW CFT. This symmetry ensures that the whole Fock space of the theory can be decomposed into multiplets which are irreducible representations of $Y(sl_N)$. In the $N = 2$ case this construction is extensively developed and it was proven that the excitations created by the so called spinon fields span the whole chiral Fock space of the theory. The excitations created by these spinon fields have shown up to behave as semionic quasiparticles.

We show that the holomorphic factors that we present for the spin $1/2$ primary field in the $SU(2)_1$ CFT exactly behave as the spinon fields of Refs. [7, 8], thus providing a fermionic

Lagrangian formulation for spinons. In this respect, our construction should be compared with the original bosonic formulation, where a heuristic holomorphic factorization of the WZW fields is assumed and no Lagrangian is given, but the dynamics is instead recovered from the conformal properties of the fields.

This letter is organized as follows: in section *ii*) we prove the holomorphic factorization of the fundamental representation $SU(N)_1$ primary fields in terms of gauge invariant fermions. In section *iii*) we discuss the fermionic (Lagrangian) representation of $SU(2)_1$ spinon fields. In section *iv*) we present the factorization of the rest of the primaries in $SU(N)_1$ and comment on their relation with the Yangian symmetry. Finally, we summarize our results in section *v*). For the sake of conciseness we also recall the fundamentals of the Yangian algebra in the appendix.

ii) Holomorphic factorization of WZW fields

It is known that the $SU(N)_1$ CFT can be formulated as a constrained fermionic model, that is as a $U(N)/U(1)$ fermionic coset theory [10]. The constraint is imposed on a system of N free Dirac fermions by requiring that physical states $|phys\rangle$ are singlet under the $U(1)$ current,

$$J_\mu|phys\rangle = 0. \quad (1)$$

This is achieved in the path integral formulation by introducing a Lagrange multiplier a_μ which acts as a $U(1)$ gauge field with no dynamics. The partition function for the constrained model in Euclidean space reads

$$\mathcal{Z} = \int \mathcal{D}\psi^\dagger \mathcal{D}\psi \mathcal{D}a_\mu \exp(-\int \mathcal{L} d^2x), \quad (2)$$

where the fermionic Lagrangian $\mathcal{L}$ is given by

$$\mathcal{L} = \frac{1}{\sqrt{2\pi}} \psi^{\dagger i} (\not{\partial} + i \not{a}) \psi^i. \quad (3)$$

Here ψ^i , $i = 1, \dots, N$, are Dirac fermions and a_μ is the abelian gauge field implementing the constraint. This procedure removes the charge degree of freedom from the fermionic system.

The theory described by the Lagrangian (3) has a local $U(1)$ invariance, thus its physical operators will be those invariant under $U(1)$ gauge transformations. In the present case they are constructed as [11]

$$\begin{aligned} \hat{\psi}^i &= \exp(-i \int_x^\infty a_\mu dx_\mu) \psi^i, \\ \hat{\psi}^{i\dagger} &= \psi^{i\dagger} \exp(i \int_x^\infty a_\mu dx_\mu). \end{aligned} \quad (4)$$

These fields are independent of the path chosen for the integration in the exponential due to the a_μ equations of motion.

It is convenient now to pass to complex coordinates $z = \frac{1}{\sqrt{2}}(x_0 + ix_1)$, $\bar{z} = \frac{1}{\sqrt{2}}(x_0 - ix_1)$ and to write the partition function in a decoupled form. This is achieved through the following

change of variables

$$\begin{aligned} a &= i(\bar{\partial}u)u^{-1} & \bar{a} &= i(\partial\bar{u})\bar{u}^{-1}, \\ \psi_1 &= u\chi_1 & \psi_2^\dagger &= \chi_2^\dagger u^{-1}, \\ \psi_2 &= \bar{u}\chi_2 & \psi_1^\dagger &= \chi_1^\dagger \bar{u}^{-1}. \end{aligned} \quad (5)$$

where $\partial \equiv \frac{\partial}{\partial z}$, $\bar{\partial} \equiv \frac{\partial}{\partial \bar{z}}$ and $\psi = \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$. The fields u and $\bar{u}$ are parametrized in terms of real scalar fields as $u = \exp(-\phi - i\eta)$ and $\bar{u} = \exp(\phi - i\eta)$.

Taking into account the gauge fixing procedure and the Jacobians associated to (5) [12] one arrives at the desired decoupled form for the partition function (2):

$$\mathcal{Z} = \mathcal{Z}_{ff} \mathcal{Z}_{fb} \mathcal{Z}_{gh}, \quad (6)$$

where

$$\begin{aligned} \mathcal{Z}_{ff} &= \int \mathcal{D}\chi^\dagger \mathcal{D}\chi \exp\left(-\frac{1}{\pi} \int (\chi_2^\dagger \bar{\partial}\chi_1 + \chi_1^\dagger \partial\chi_2) d^2x\right), \\ \mathcal{Z}_{fb} &= \int \mathcal{D}\phi \exp\left(\frac{N}{2\pi} \int \phi \Delta \phi d^2x\right), \\ \mathcal{Z}_{gh} &= \int \mathcal{D}\bar{c} \mathcal{D}c \mathcal{D}\bar{b} \mathcal{D}b \exp\left(-\int (b\partial c + \bar{b}\bar{\partial}\bar{c}) d^2x\right). \end{aligned} \quad (7)$$

Notice that, although the partition function of the theory is completely decoupled, BRST quantization conditions connect the different sectors in order to ensure unitarity [13]. Once we have obtained the desired decoupled expression for the partition function of the $SU(N)_1$ model we proceed further to study its spectrum.

In the decoupled picture, the components of the gauge invariant fields (4) can be written as

$$\begin{aligned} \hat{\psi}_1^i(z) &= e^{-\varphi(z)} \chi_1^i(z) & \hat{\psi}_1^{i\dagger}(\bar{z}) &= \chi_1^{i\dagger}(\bar{z}) e^{-\bar{\varphi}(\bar{z})} \\ \hat{\psi}_2^i(\bar{z}) &= e^{\bar{\varphi}(\bar{z})} \chi_2^i(\bar{z}) & \hat{\psi}_2^{i\dagger}(z) &= \chi_2^{i\dagger}(z) e^{\varphi(z)} \end{aligned} \quad (8)$$

where

$$\begin{aligned} \varphi(z) &= \phi + i \int_x^\infty dz_\mu \epsilon_{\mu\nu} \partial_\nu \phi \\ \bar{\varphi}(\bar{z}) &= \phi - i \int_x^\infty dz_\mu \epsilon_{\mu\nu} \partial_\nu \phi \end{aligned} \quad (9)$$

are the chiral (holomorphic and anti-holomorphic) components of the free boson ϕ , satisfying

$$\bar{\partial}\varphi = \partial\bar{\varphi} = 0. \quad (10)$$

This fact together with the equation of motion of the free fermions χ ensures that $\hat{\psi}_1^i$ and $\hat{\psi}_2^{i\dagger}$ ($\hat{\psi}_2^i$ and $\hat{\psi}_1^{i\dagger}$) are holomorphic (anti-holomorphic).

Eq.(8) makes evident the relevance of the gauge invariance requirement for physical operators: physical excitations created by gauge invariant composites $\hat{\psi}_1^i$, $\hat{\psi}_2^{i\dagger}$ are automatically holomorphic, while those created by $\hat{\psi}_2^i$, $\hat{\psi}_1^{i\dagger}$ are anti-holomorphic.

Since the composite fermions in (4) are now completely given in terms of free fields, their properties can be easily studied. In particular, we are interested in their short distance

operator product expansion (OPE). Being the composite fermions primary fields, the leading singularity in their OPE is governed by their conformal dimensions h and $\bar{h}$. This in turn is simply given by the sum of the dimensions of their decoupled constituents (see eqs.(8) and (9)). For the holomorphic composites $\hat{\psi}_1^i$ and $\hat{\psi}_2^{i\dagger}$ one has

$$h = \frac{1}{2} - \frac{1}{2N} = \frac{N-1}{2N}, \quad \bar{h} = 0. \quad (11)$$

(Similarly, $h = 0$ and $\bar{h} = \frac{N-1}{2N}$ for the anti-holomorphic composites $\hat{\psi}_2^i$ and $\hat{\psi}_1^{i\dagger}$).

This altogether gives place to the following OPE:

$$\hat{\psi}_2^{i\dagger}(z)\hat{\psi}_1^j(w) = \frac{\delta^{ij}}{(z-w)^{(N-1)/N}} + \dots \quad (12)$$

(Analogously $\hat{\psi}_1^{i\dagger}(\bar{z})\hat{\psi}_2^j(\bar{w}) = \frac{\delta^{ij}}{(\bar{z}-\bar{w})^{(N-1)/N}} + \dots$).

These composite fields can indeed be seen as the holomorphic factors of the WZW primaries $g, g^\dagger$ in the fundamental representation of $SU(N)$. This fact follows from the consideration of the bosonization dictionary [10] and the non observability of phase factors in fermion bilinears, which allow us to write

$$g^{ij}(z, \bar{z}) = \psi_2^i \psi_2^{j\dagger} = \hat{\psi}_2^i(\bar{z})\hat{\psi}_2^{j\dagger}(z) \quad (13)$$

This equation provides the explicit holomorphic factorization of the WZW primary field g . In the same way one gets

$$(g^\dagger)^{ij}(z, \bar{z}) = \psi_1^i \psi_1^{j\dagger} = \hat{\psi}_1^i(z)\hat{\psi}_1^{j\dagger}(\bar{z}). \quad (14)$$

Eqs. (13) and (14) define the holomorphic factors of g and $g^\dagger$ up to an arbitrary constant phase.

Let us recall that the factors $\hat{\psi}$ and $\hat{\psi}^\dagger$ independently create physical excitations in contrast with the original fermions which do not. This result is one of the main points in our work, that is the explicit factorization of the g WZW primary field in terms of holomorphic and anti holomorphic gauge invariant fermions.

iii) $SU(2)_1$ spinons

We will now show that the holomorphic gauge-invariant composite fermions in the fermionic coset formulation of the $SU(2)_1$ CFT do fulfill the characterization of the holomorphic part of the spin 1/2 primary field. In particular, we show that they provide a Lagrangian construction for spinon fields.

The spinon formulation of $SU(2)_1$ WZW theory has been recently developed [7, 8]. It exploits the representation theory of the Yangian $Y(sl_2)$ algebra (see the appendix), a symmetry present in the model, in order to give a new representation of its chiral spectrum. The spectrum thus obtained is naturally organized into Yangian multiplets, each one associated

to a Yangian highest weight state, and turns out to be exactly equivalent to the standard construction based on the representation theory of Virasoro and Kac-Moody algebras.

One can construct the Yangian highest weight states in the context the $SU(2)_1$ WZW theory [6, 8]. Moreover, in these references each state in the spinon Fock space is constructed by the repeated action of the Laurent modes of a spinon field on the vacuum (we summarize this construction below), where the spinon field is defined as the “holomorphic part” of the spin 1/2 Virasoro primary field. Rigorously speaking, one assumes a heuristic holomorphic factorization of the WZW primary fields $g(z, \bar{z}) = \phi(z)\bar{\phi}(\bar{z})$ and then makes use of OPEs that follow from the action of Virasoro and Kac-Moody generators on primary fields, as was done in Ref. [8]. An alternative construction was given in Ref. [7] in terms of chiral vertex operators motivated by the bosonization of two complex Dirac fermions with an $SU(2)$ charge degree of freedom arbitrarily wiped out.

In our construction below the holomorphic factors directly follow from the fermionic coset Lagrangian (3) and from the principle of BRST invariance for physical operators. Also their OPEs can be derived from the partition function by standard field theoretical methods. The charge degree of freedom of the original fermions is also naturally removed from the theory by the coset constraining procedure, in a situation which is reminiscent of what happens in the original scenario for spinons, namely the Haldane-Shastry spin chain, where the charge degree of freedom is fixed by construction [14, 15].

In order to make contact with the spinon formulation discussed in Ref.[8] we are going to first study some details of the $SU(2)_1$ case. In particular, the $SU(2)$ fundamental representation and its conjugate are equivalent, so the matrix elements of the WZW field g and its dagger $g^\dagger$ are related by

$$g^{ij} = -\epsilon^{jl}(g^\dagger)^{lk}\epsilon^{ki}. \quad (15)$$

This relation does not trivially apply to the fermi bilinears (13) and (14), since it leads to inconsistencies in correlators. As an example, consider the two-point correlator [16, 17]

$$\langle g^{ij}(g^\dagger)^{kl} \rangle = -\epsilon^{kr}\langle g^{ij}g^{rs} \rangle \epsilon^{sl} \neq 0. \quad (16)$$

The inconsistency is apparent when one uses the bosonization rules to evaluate $\langle g^{ij}g^{rs} \rangle$ in terms of fermions, thus obtaining a vanishing result. Notice that a similar situation is already present in Witten’s original bosonization rules for the $U(N)$ case [18].

To circumvent this problem one can use the bosonization rules with the proviso that the fields g and $g^\dagger$ must appear pairwised. In other words, an expression like $g^{ij}g^{rs}$ must be written as $-\epsilon^{jl}(g^\dagger)^{lk}\epsilon^{ki}g^{rs}$ *before* using the rules (13)-(14). We can write down the same rule in terms of gauge invariant fermions by stating that the substitutions

$$g^{ij}(z, \bar{z}) \leftrightarrow \begin{cases} \hat{\psi}_2^i(\bar{z})\hat{\psi}_2^{j\dagger}(z) & \text{when } g \text{ occurs as an odd factor} \\ \epsilon^{jl}\hat{\psi}_1^l(z)\epsilon^{ik}\hat{\psi}_1^{k\dagger}(\bar{z}) & \text{when } g \text{ occurs as an even factor} \end{cases} \quad (17)$$

are the appropriate ones when dealing with a multi-point product of g -fields. In this respect, the literature about the spinon formulation of $SU(2)_1$ WZW theory [8] makes reference to the holomorphic part of the g -field only and makes implicit use of the relation (15) any time that

a spinon field occurs as an even factor in a multi-spinon operator product. This gives rise to two different “sectors” in the spinon Fock space: the vacuum sector, consisting of states created by an even number of spinon operators, and the spin 1/2 sector, containing states created by an odd number of such operators. To conform to the usual notation, we then define a holomorphic field $\sigma^j(z)$ as:

$$\sigma^j(z) = \begin{cases} \hat{\psi}_2^{j\dagger}(z) & \text{when acting on the vacuum sector} \\ \epsilon^{jk} \hat{\psi}_1^k(z) & \text{when acting on the spin 1/2 sector} \end{cases} \quad (18)$$

(The phase for each holomorphic factor is chosen for later convenience; it is of course irrelevant while considering g correlators).

The σ - σ OPE is now readily evaluated by using eq.(12). It is given by

$$\sigma^i(z)\sigma^j(w) = (-1)^q \frac{\epsilon^{ij}}{(z-w)^{1/2}} + \dots \quad (19)$$

where $q = 0$ when this product acts on the vacuum sector and $q = 1$ otherwise. As pointed out in [8], it is this OPE what characterizes the σ 's as spinon fields. We have then shown that the holomorphic gauge-invariant composite fermions, as given in the σ^j definition, provide a Lagrangian construction for spinon fields.

In view of the discussion above, the σ^j field fulfills the properties of the bosonic spinon field. Once this identification is made, one can follow the same procedure described in [7, 8] to construct the full chiral Hilbert space of the $SU(2)_1$ CFT from the action of the modes of the σ field. For the sake of conciseness we summarize this construction here.

Due to the square root branch cut in the OPE singularity (19) the excitations created by the spinon fields are semionic. This in turn implies that their mode expansion will involve integer (half-integer) powers of z when acting on the vacuum (spin 1/2) sector:

$$\sigma^i(z) = \sum_{m=0}^{\infty} z^{m+\frac{q}{2}} \sigma_{-m-\frac{1}{4}-\frac{q}{2}}^i, \quad (20)$$

where

$$\sigma_{-m+\frac{3}{4}-\frac{q}{2}}^i = \oint \frac{dz}{2\pi i} z^{-m-\frac{q}{2}} \sigma^i(z). \quad (21)$$

These modes satisfy generalized commutation relations which can be obtained from (24), giving

$$\sum_{l=0}^{\infty} C_l^{(-1/2)} \left(\sigma_{-m-\frac{q+1}{2}-l+\frac{3}{4}}^i \sigma_{-n-\frac{q}{2}+l+\frac{3}{4}}^j - \sigma_{-n-\frac{q+1}{2}-l+\frac{3}{4}}^j \sigma_{-m-\frac{q}{2}+l+\frac{3}{4}}^i \right) = (-1)^q \epsilon^{ij} \delta_{m+n+q-1,0} \quad (22)$$

where $C_l^{(-1/2)}$ are the coefficients of the Taylor expansion of $(1-z)^{-1/2}$ [17].

The chiral part of the Fock space of the $SU(2)_1$ conformal field theory can be constructed in terms of these modes. As stated above, this space can be classified into multiplets corresponding to the irreducible representations of the Yangian algebra $Y(sl_2)$. Each multiplet is

constructed from a highest weight state by the action of the Yangian generators. The highest weight states can be constructed in terms of the spinon modes in the following way: one has to take linear combinations of M -spinon states

$$\sigma_{-n_1-1/4}^1 \cdots \sigma_{-n_M-1/4}^1 |0\rangle \quad (23)$$

that are annihilated by the raising operators Q_0^+ and Q_1^+ . These states correspond to the “fully polarized M -spinon states” of Haldane [2]. The defining level 0 and 1 generators of $Y(sl_2)$ can be written in terms of the modes of the Kac-Moody currents [6] as

$$\begin{aligned} Q_0^a &= J_0^a, \\ Q_1^a &= \frac{1}{2} f_{bc}^a \sum_{m=0}^{\infty} : J_{-m}^b J_m^c :. \end{aligned} \quad (24)$$

The irreducible Yangian multiplet corresponding to a given highest weight state is then constructed by repeatedly applying the generators Q_0^- and Q_1^- . Finally, the union of all of these multiplets forms a basis of the full Hilbert space.

iv) Holomorphic factorization of other primaries in $SU(N)_1$ and their relation with $SU(N)_1$ spinon fields

In section *ii)* we presented the holomorphic factorization of the primary field g in the $SU(N)_1$ WZW theory. In the general case $N > 2$ the theory contains N primary fields, each one corresponding to an integrable representation, which can be constructed as suitable symmetrized products of fields in the fundamental representation. We will now show how the holomorphic factors of all these primaries can be explicitly constructed in terms of the gauge invariant fermions (8).

The integrable representations for the $SU(N)_1$ case correspond to Young diagrams with one column and at most $N - 1$ rows. The primary fields in these representations, Φ_p , are then obtained by taking normal-ordered antisymmetric products of p fields ($p = 0, 1, \dots, N - 1$) in the fundamental representation [10].

In terms of the composite fermions (8) the holomorphic parts of the primaries corresponding to $p = 2, \dots, N - 1$ are given by

$$\Phi_p^{i_1, i_2, \dots, i_p}(z) = \mathcal{A} \left(: \hat{\psi}_1^{i_1} \hat{\psi}_1^{i_2} \dots \hat{\psi}_1^{i_p} : \right), \quad (25)$$

while in terms of the decoupled fields they can be written as

$$\Phi_p^{i_1, i_2, \dots, i_p}(z) = \mathcal{A} \left(: \chi_1^{i_1} \chi_1^{i_2} \dots \chi_1^{i_p} : \right) : \exp(-p\varphi) :. \quad (26)$$

The conformal dimensions of these fields can be evaluated as the sum of two independent contributions: $p/2$ coming from the p free fermions and $-p^2/2N$ coming from the vertex operator $: \exp(-p\varphi) :$. The result is

$$h_p = \frac{p}{2} - \frac{p^2}{2N} = \frac{p(N-p)}{2N}. \quad (27)$$

These dimensions agree with the conformal dimensions of the $SU(N)_1$ primary fields (see for instance [16]), each one in the representation Λ_p (associated with a Young diagram with p vertical boxes), given by

$$h_{\Lambda_p} = \frac{c_{\Lambda_p}}{c_v + k} \quad (28)$$

where $c_v = N$ for $SU(N)$, $k = 1$ is the Kac-Moody level and $c_{\Lambda_p} = \frac{p}{2N}(N+1)(N-p)$ is the Casimir of the representation Λ_p .

The Yangian structure is also present in the $N > 2$ case [9], but the spinon construction has not yet been presented.

v) Conclusions

We have proved that the physical operators in the fermionic coset realization of $SU(N)_1$ CFT are automatically holomorphic or antiholomorphic. The standard bosonization rules, when written in terms of gauge invariant fermions, provide an explicit holomorphic factorization of WZW primary fields. Moreover, we have shown how to express the holomorphic part of any primary in the $SU(N)_1$ theory.

As expected, the holomorphic part of the spin 1/2 primary field in the $SU(2)_1$ case fulfills the characterization of the so called spinon field. As a consequence, the fermionic coset model $U(2)/U(1)$ provides a Lagrangian formulation for spinons, which are known to behave as semionic 1 + 1 dimensional quasiparticles.

The present description of the spinon representation of $SU(2)$ CFT in terms of composite fermions in a constrained fermionic formulation seems to be, in our opinion, a natural framework to investigate some open questions. In the same line, the fundamental rôle played by composite gauge invariant fermions has already shown up in the coset formulation of the Ising model [11] (see also [19]), where they were used to build up the order and disorder operators. We expect that this guideline should allow for the holomorphic factorization of some primaries in more general coset models.

It will be also interesting to see whether an underlying structure similar to the Yangian appearing in $SU(N)_1$ is also present in other CFTs, such as higher level $SU(N)$ CFT or coset models. In fact, the holomorphic factorization of the WZW primaries in the $SU(N)_k$ case ($k > 1$) in terms of gauge invariant fermions can be envisaged following the same lines as those presented here. The connection with the spinon basis constructed in Ref. [20] will be discussed elsewhere [21].

Acknowledgements: we would like to thank J. Fuchs and K. Schoutens for useful discussions and specially E.F.Moreno for his helpful suggestions and valuable comments. We also thank F. A. Schaposnik for comments on the preliminar manuscript.

Note added: after this paper was finished, the spinon construction has been extended to the $N > 2$ case [24]. The complete (chiral) Fock space has been constructed there by acting on the vacuum with the modes of the holomorphic factor of the primary field transforming in the $\bar{N}$ representation of $SU(N)$ (in their notation, $\phi^\alpha(z)$, $\alpha = 1, 2, \dots, N$). As stressed in [24], the statistics of these fields is encoded in their OPE. In their notation, it is given by

$$\phi(z)\phi(w) = \frac{\phi'(w)}{(z-w)^{1/N}} + \dots \quad (29)$$

where ϕ' is the primary field in the representation Λ_{N-2} of $SU(N)$.

This construction can of course be made out from the fields we present in section *iv*). In particular, it is simpler in our notation to work with the representation N instead of $\bar{N}$, and define the spinon creation operators from the modes of $\Phi_{p=1}$. Using our expression (26) for this field, we can easily evaluate the OPE in terms of the decoupled fields:

$$\Phi_1^i(z)\Phi_1^j(w) = \frac{1}{(z-w)^{1/N}} \mathcal{A}(:\chi_1^i(w)\chi_1^j(w):) : \exp(-2\varphi(w)) : + \dots = \frac{1}{(z-w)^{1/N}} \Phi_2^{ij}(w) + \dots \quad (30)$$

The agreement between (29) and (30) shows that the gauge invariant fermions (4) provide an explicit Lagrangian realization of the spinon fields not only for $N = 2$ but for arbitrary N .

Appendix: Yangian structure of the Fock space of $SU(N)_1$ conformal field theories

We review here the main aspects of a novel representation of the spectrum of the chiral part of $SU(N)_1$ CFTs, first introduced by Haldane et al.[2, 6] and further developed in [7, 8, 9, 20]. It was originally presented in [2] as an extrapolation of the exact solution of long range interaction 1D spin chains [14, 15] and the fact that this chain is described in the low energy limit by the level one $SU(2)$ conformal field theory [22].

In the $SU(2)_1$ case the new representation, known as spinon representation, allows for a quasi-particle interpretation: in this sense, each state corresponds to a given number of spin one-half, neutral, semionic, free quasi-particles (spinons).

The quantum numbers that characterize the representation correspond to the behavior of the states under the action of the generators Q_n^a of an infinite algebra, known as Yangian algebra [23] (where $n = 0, 1, \dots$ and a is a Lie algebra index). The Yangian $Y(\hat{g})$ associated to a Lie algebra $\hat{g}$ is a Hopf algebra that is neither commutative nor cocommutative, being a simple non-trivial example of a quantum group.

The Yangian was defined in [23] by starting with the generators Q_0^a and Q_1^a , which for $\hat{g} = sl_N$ (the Lie algebra of $SU(N)$) are defined by the relations

$$\begin{aligned} [Q_0^a, Q_0^b] &= f^{abc} Q_0^c, \\ [Q_0^a, Q_1^b] &= f^{abc} Q_1^c, \\ [Q_1^a, [Q_1^b, Q_0^c]] + (\text{cyclic in } a, b, c) &= A^{abcdef} \{Q_0^d, Q_0^e, Q_0^f\} \\ [[Q_1^a, Q_1^b], [Q_0^c, Q_1^d]] + [[Q_1^c, Q_1^d], [Q_0^a, Q_1^b]] &= (A^{abpqrs} f^{cdp} + A^{cdpqrs} f^{abp}) \{Q_0^q, Q_0^r, Q_1^s\}, \end{aligned} \quad (31)$$

where $A^{abpdef} = \frac{1}{4} f^{adp} f^{beq} f^{cfr} f^{pqr}$ and $\{..., ..., ...\}$ denotes a completely symmetrized product. Higher level generators are recursively defined for $n \geq 2$ by

$$c_v Q_n^a = f^{abc} [Q_1^c, Q_{n-1}^b], \quad (32)$$

where c_v is the Casimir in the adjoint representation.

A central property of this algebra is the comultiplication rule that can be viewed as a generalization of angular momentum addition:

$$\begin{aligned} \Delta_{\pm}(Q_0^a) &= Q_0^a \otimes 1 + 1 \otimes Q_0^a \\ \Delta_{\pm}(Q_1^a) &= Q_1^a \otimes 1 + 1 \otimes Q_1^a \pm \frac{1}{2} f^{abc} Q_0^b \otimes Q_0^c. \end{aligned} \quad (33)$$

Being the Yangian a symmetry of the $SU(N)_1$ CFT the whole Hilbert space can be decomposed into a sum of multiplets corresponding to its irreducible representations. Each multiplet is characterized by a highest weight state defined by

$$Q_n^{\alpha} |\psi\rangle = 0 \quad (34)$$

where α corresponds to a positive root in the Cartan basis. Its vectors are eigenstates of Q_n^i , where i corresponds to Cartan subalgebra generators. Each highest weight state is characterized by a finite set of non-decreasing integers $\{n_i\}_{i=1, \dots, p}$. In the $SU(2)_1$ quasi-particle language each one of these integers represents one spinon in the n_i -th orbital.

References

- [1] See e.g. “*Fractional Statistics and anyon superconductivity*”, ed. F. Wilczek, World Scientific (Singapore), 1989.
- [2] F.D.M.Haldane, Phys. Rev. Lett. **66**, 1529 (1991).
- [3] F.D.M.Haldane, Phys. Rev. Lett. **67**, 937 (1991).
- [4] Y.S.Wu, Phys. Rev. Lett. **73**, 922 (1994).
- [5] X-G. Wen, Y-S. Wu and Y. Hatsugai, Nucl. Phys. **B422**, 476 (1994).
- [6] F.D.M.Haldane, Z.N.C.Ha, J.C.Talstra, D.Bernard and V.Pasquier, Phys. Rev. Lett. **69**, 2021 (1992).
- [7] D.Bernard, V.Pasquier and D.Serban, Nucl. Phys. **B428** [FS], 612 (1994).
- [8] P.Bouwknegt, A.W.W.Ludwig and K.Schoutens, Phys. Lett. **338B**, 448 (1994).
- [9] K.Schoutens, Phys. Lett. **331B**, 335 (1994).
- [10] S.G.Naculich and H.J.Schnitzer, Nucl. Phys. **B332**, 583, (1990).
- [11] D.C.Cabra and K.Rothe, Phys. Rev. **D51** R2509 (1995).
- [12] A.Polyakov and P.Wiegmann, Phys.Lett. **131B**, 121 (1983); R.E.Gamboa Saraví F.A.Schaposnik and J.E.Solomin, Nucl.Phys.**B185**, 238 (1981).
- [13] D.Karabali, Q.Park, H.Schnitzer and Z.Yang, Phys. Lett. **B216**, 307, (1989).
- [14] F.D.M.Haldane, Phys. Rev. Lett. **60**, 635 (1988).
- [15] B.S.Shastry, Phys. Rev. Lett. **60**, 639 (1988).
- [16] V.G.Knizhnik and A.B.Zamolodchikov, Nucl. Phys. **B247**, 83 (1984).
- [17] A.B.Zamolodchikov and V.A.Fateev, Sov. Phys. JETP **62**, 215 (1985).
- [18] E.Witten, Commun. Math. Phys. **92**, 455 (1984).
- [19] D.C.Cabra and E.F.Moreno, Nucl. Phys. **B475**, 522 (1996).
- [20] P.Bouwknegt, A.W.W.Ludwig and K.Schoutens, Phys. Lett. **359B**, 304 (1995).
- [21] D.C.Cabra, E.F.Moreno and G.L.Rossini, *Quasiparticles in $SU(2)_k$ fermion cosets*, unpublished.
- [22] I.Affleck, Phys. Rev. Lett. **55**, 1355 (1985).

- [23] V.G.Drinfel'd, Sov. Math. Dokl. **32**, 254 (1985).
- [24] P.Bouwknegt and K.Schoutens, *The $SU(n)_1$ WZW models, spinon decomposition and yangian structure*, hep-th/9607064.